

# Second Order Periodic Problem with $\phi$ -Laplacian and Impulses

Irena Rachůnková\* and Milan Tvrdý†

March 30, 2004

**Abstract.** Existence principles for the BVP  $(\phi(u'))' = f(t, u, u')$ ,  $u(t_i+) = J_i(u(t_i))$ ,  $u'(t_i+) = M_i(u'(t_i))$ ,  $i = 1, 2, \dots, m$ ,  $u(0) = u(T)$ ,  $u'(0) = u'(T)$  are presented. They are based on the method of lower/upper functions which need not be well-ordered.

**Mathematics Subject Classification 2000.** 34B37, 34B15, 34C25

**Keywords.**  $\phi$ -Laplacian, impulses, lower/upper functions, periodic solutions

## 1. Formulation of the problem

Let  $m \in \mathbb{N}$ ,  $0 = t_0 < t_1 < \dots < t_m < t_{m+1} = T$  and  $D = \{t_1, t_2, \dots, t_m\}$ . Define  $\mathbb{C}_D$  (or  $\mathbb{C}_D^1$ ) as the sets of functions  $u : [0, T] \mapsto \mathbb{R}$ ,

$$u(t) = \begin{cases} u_{[0]}(t) & \text{if } t \in [0, t_1], \\ u_{[1]}(t) & \text{if } t \in (t_1, t_2], \\ \dots & \dots \\ u_{[m]}(t) & \text{if } t \in (t_m, T], \end{cases}$$

where  $u_{[i]}$  is continuous on  $[t_i, t_{i+1}]$  (or continuously differentiable on  $[t_i, t_{i+1}]$ ) for  $i = 0, 1, \dots, m$ . We put  $\|u\|_D = \|u\|_\infty + \|u'\|_\infty$ , where  $\|u\|_\infty = \sup \operatorname{ess}_{t \in [0, T]} |u(t)|$ . Then  $\mathbb{C}_D$  and  $\mathbb{C}_D^1$  respectively with the norms  $\|\cdot\|_\infty$  and  $\|\cdot\|_D$  become Banach spaces. Further,  $\mathbb{AC}_D$  is the set of functions  $u \in \mathbb{C}_D$  which are absolutely continuous on each subinterval  $(t_i, t_{i+1})$ ,  $i = 0, 1, \dots, m$ .

We consider the problem

$$(1.1) \quad (\phi(u'(t)))' = f(t, u(t), u'(t)) \quad \text{a.e. on } [0, T],$$

$$(1.2) \quad u(t_i+) = J_i(u(t_i)), \quad u'(t_i+) = M_i(u'(t_i)), \quad i = 1, 2, \dots, m,$$

$$(1.3) \quad u(0) = u(T), \quad u'(0) = u'(T),$$

---

\*Supported by the grant No. 201/04/1077 of the Grant Agency of the Czech Republic and by the Council of Czech Government J14/98:153100011

†Supported by the grant No. 201/04/1077 of the Grant Agency of the Czech Republic

where  $u'(t_i) = u'(t_i-) = \lim_{t \rightarrow t_i-} u'(t)$  for  $i = 1, 2, \dots, m+1$ ,  $u'(0) = u'(0+) = \lim_{t \rightarrow 0+} u'(t)$ ,  $f$  is an  $\mathbb{L}_1$ -Carathéodory function, functions  $J_i, M_i$  are continuous on  $\mathbb{R}$  and  $\phi$  is an increasing homeomorphism such that  $\phi(0) = 0$  and  $\phi(\mathbb{R}) = \mathbb{R}$ . A typical example of a proper function  $\phi$  is the  $p$ -Laplacian  $\phi_p(y) = |y|^{p-2}y$ , where  $p > 1$ .

A *solution* of the problem (1.1)–(1.3) is a function  $u \in \mathbb{C}_D^1$  such that  $\phi(u') \in \mathbb{AC}_D$  and (1.1)–(1.3) hold.

A function  $\sigma \in \mathbb{C}_D^1$  is called a *lower function* of (1.1)–(1.3) if  $\phi(\sigma') \in \mathbb{AC}_D$  and

$$(1.4) \quad \begin{cases} \phi(\sigma'(t))' \geq f(t, \sigma(t), \sigma'(t)) & \text{for a.e. } t \in [0, T], \\ \sigma(t_i+) = J_i(\sigma(t_i)), \sigma'(t_i+) \geq M_i(\sigma'(t_i)), & i = 1, 2, \dots, m, \\ \sigma(0) = \sigma(T), \sigma'(0) \geq \sigma'(T). \end{cases}$$

Similarly, a function  $\sigma \in \mathbb{C}_D^1$  with  $\phi(\sigma') \in \mathbb{AC}_D$  is an *upper function* of (1.1)–(1.3) if it satisfies the relations (1.4) but with reversed inequalities.

The aim of this paper is to offer existence principles for problem (1.1)–(1.3) in terms of lower/upper functions. Hence our basic assumption is the existence of lower/upper functions. We will suppose that either

$$(1.5) \quad \begin{aligned} &\sigma_1 \quad \text{and} \quad \sigma_2 \quad \text{are respectively lower and upper functions of (1.1)–(1.3)} \\ &\text{such that } \sigma_1 \leq \sigma_2 \quad \text{on } [0, T] \end{aligned}$$

or

$$(1.6) \quad \begin{aligned} &\sigma_1 \quad \text{and} \quad \sigma_2 \quad \text{are respectively lower and upper functions of (1.1)–(1.3)} \\ &\text{such that } \sigma_1 \not\leq \sigma_2 \quad \text{on } [0, T], \text{ i.e. } \sigma_1(\tau) > \sigma_2(\tau) \text{ for some } \tau \in [0, T]. \end{aligned}$$

Note that problems with  $\phi$ -Laplacians and impulses have not been studied yet. As concerns problem (1.1), (1.3) (without impulses), there are various results about its solvability. For example the papers [4] and [19] present some results about the existence or multiplicity of periodic solutions of the equation

$$(1.7) \quad (\phi_p(u'))' = f(t, u)$$

under non resonance conditions imposed on  $f$ . The paper [10] presents general existence principles for the vector problem (1.1), (1.3). Using this the authors provide various existence theorems and illustrative examples. The vector case is also considered in [9], [11] and [12]. The existence of periodic solutions of the Liénard type equations with  $p$ -Laplacians has been proved in the resonance case under the Landesman-Lazer conditions in [5] and [6]. Multiplicity results of the Ambrosetti-Prodi type for this problem (with a real parameter) can be found in [8].

The papers which are devoted to the lower/upper functions method for the problem (1.1), (1.3) mostly deal with the condition (1.5), i.e. they assume well-ordered

$\sigma_1/\sigma_2$ . We can refer to the papers [1] and [3] which study the problem (1.1), (1.3) under the Nagumo type two-sided growth conditions and to the paper [17] where the second order equation with a  $\phi$ -Laplacian is considered provided a functional right-hand side of this equation fulfils one-sided growth conditions of the Nagumo type. The significance of the lower/upper functions method is shown in the papers [7] and [18] where this method is used in the investigation of singular periodic problems with a  $\phi$ -Laplacian. The paper [2] is, to our knowledge, the only one presenting the lower/upper functions method for the problem (1.7), (1.3) (with a  $\phi$ -Laplacian) under the assumption that  $\sigma_1 \geq \sigma_2$ , i.e. lower/upper functions are in the reverse order. If  $\phi = \phi_p$  the authors get the solvability of (1.7), (1.3) for  $1 < p \leq 2$ , only. Therefore the existence principles (Theorems 2.3 and 2.4) which we state here for the impulsive problem (1.1)–(1.3) and the case (1.6) are new even for the non-impulsive problem (1.1), (1.3).

We will work with the following assumptions, where the sets  $A_i, B(t) \subset \mathbb{R}$ ,  $t \in [0, T]$ , will be determined later, according to whether (1.5) or (1.6) is assumed:

$$(1.8) \quad \begin{cases} x > \sigma_1(t_i) \implies J_i(x) > J_i(\sigma_1(t_i)) \\ x < \sigma_2(t_i) \implies J_i(x) < J_i(\sigma_2(t_i)) \end{cases} \quad \text{for } x \in A_i, \quad i = 1, 2, \dots, m;$$

$$(1.9) \quad \begin{cases} y \leq \sigma'_1(t_i) \implies M_i(y) \leq M_i(\sigma'_1(t_i)), \\ y \geq \sigma'_2(t_i) \implies M_i(y) \geq M_i(\sigma'_2(t_i)), \end{cases} \quad i = 1, 2, \dots, m;$$

$$(1.10) \quad \begin{cases} \text{there is } h \in \mathbb{L}_1 \text{ such that} \\ |f(t, x, y)| \leq h(t) \text{ for a.e. } t \in [0, T] \text{ and all } x, y \in \mathbb{R}; \end{cases}$$

$$(1.11) \quad \begin{cases} \text{there are } \omega : [0, \infty) \mapsto (0, \infty) \text{ continuous and } h \in \mathbb{L}_1 \text{ such that} \\ \int_0^\infty \frac{ds}{\omega(s)} = \infty \text{ and } |f(t, x, y)| \leq \omega(\phi(|y|)) (|y| + h(t)) \\ \text{for a.e. } t \in [0, T], \text{ all } x \in B(t) \text{ and } |y| \geq 1, \end{cases}$$

$$(1.12) \quad \begin{cases} \text{there are } c_j, d_j \in \mathbb{R}, \quad c_j \leq \sigma'_k(t) \leq d_j \text{ on } (t_{j-1}, t_j], \quad k = 1, 2, \\ \text{such that } f(t, x, c_j) \leq 0, \quad f(t, x, d_j) \geq 0 \text{ for a.e. } t \in (t_{j-1}, t_j] \\ \text{and all } x \in B(t), \quad j = 1, 2, \dots, m+1, \text{ and } c_1 \geq c_{m+1}, \quad d_1 \leq d_{m+1}, \\ M_i(c_i) \leq c_{i+1}, \quad M_i(d_i) \geq d_{i+1}, \quad i = 1, 2, \dots, m. \end{cases}$$

## 2. Main results

Below we formulate our main results:

### I. EXISTENCE PRINCIPLES FOR WELL-ORDERED CASE

**2.1 Theorem.** *Assume that (1.5), (1.8) with  $A_i = [\sigma_1(t_i), \sigma_2(t_i)]$ ,  $i = 1, 2, \dots, m$ , (1.9) and (1.11) with  $B(t) = [\sigma_1(t), \sigma_2(t)]$  hold. Then the problem (1.1) – (1.3) has*

a solution  $u$  satisfying

$$(2.1) \quad \sigma_1 \leq u \leq \sigma_2 \quad \text{on } [0, T].$$

**2.2 Theorem.** Assume that (1.5), (1.8) with  $A_i = [\sigma_1(t_i), \sigma_2(t_i)]$ ,  $i = 1, 2, \dots, m$ , (1.9) and (1.12) with  $B(t) = [\sigma_1(t), \sigma_2(t)]$  hold.

Then the problem (1.1) – (1.3) has a solution  $u$  satisfying (2.1) and

$$(2.2) \quad c_j \leq u'(t) \leq d_j \quad \text{for } t \in (t_{j-1}, t_j], \quad j = 1, 2, \dots, m + 1.$$

## II. EXISTENCE PRINCIPLES FOR NON-ORDERED CASE

**2.3 Theorem.** Assume that (1.6), (1.8) with  $A_i = \mathbb{R}$ ,  $i = 1, 2, \dots, m$ , (1.9) and (1.10) hold. Then the problem (1.1) – (1.3) has a solution  $u$  satisfying

$$(2.3) \quad |u(t_u)| \leq \max\{|\sigma_1(t_u)|, |\sigma_2(t_u)|\} \quad \text{for some } t_u \in [0, T].$$

**2.4 Theorem.** Assume that (1.6), (1.8) with  $A_i = \mathbb{R}$ ,  $i = 1, 2, \dots, m$ , (1.9) and (1.12) with  $B(t) = \mathbb{R}$  hold. Then the problem (1.1) – (1.3) has a solution  $u$  satisfying (2.2) and (2.3).

Note that Theorems 2.2 and 2.4 impose no growth restrictions on  $f$ . For example, taking  $f(t, x, y) = y(y^{2k}x^{2n} + 1) - x^{2n-1} + e(t)$ , where  $e \in \mathbb{C}_D$ ,  $k, n \in \mathbb{N}$ , we can check that there are  $c_j \in (-\infty, 0)$ ,  $d_j \in (0, \infty)$ ,  $j = 1, 2, \dots, m + 1$ , such that  $c_1 \geq c_{m+1}$ ,  $d_1 \leq d_{m+1}$ ,  $f(t, x, c_j) \leq 0$  and  $f(t, x, d_j) \geq 0$  for a.e.  $t \in (t_{j-1}, t_j]$  and all  $x \in \mathbb{R}$ ,  $j = 1, 2, \dots, m + 1$ .

## 3. A fixed point operator

We will transform the problem (1.1)–(1.3) into a fixed point problem in  $\mathbb{C}_D^1$ . First, we borrow some ideas from [10] to get the following two lemmas.

**3.1 Lemma.** For each  $\ell \in \mathbb{C}_D$  and  $d \in \mathbb{R}$ , the function

$$\Psi_{\ell, d} : \mathbb{R} \mapsto \mathbb{R}, \quad \Psi_{\ell, d}(a) = d + \int_0^T \phi^{-1}(a + \ell(t)) \, dt$$

has exactly one zero point  $a(\ell, d)$  in  $\mathbb{R}$ .

*Proof.* Choose  $\ell \in \mathbb{C}_D$  and  $d \in \mathbb{R}$ . Since  $\Psi_{\ell, d}$  is continuous, increasing on  $\mathbb{R}$  and  $\Psi_{\ell, d}(\mathbb{R}) = \mathbb{R}$ , there is a unique real number  $a(\ell, d)$  such that

$$(3.1) \quad \Psi_{\ell, d}(a(\ell, d)) = 0. \quad \square$$

**3.2 Lemma.** *The mapping  $a : \mathbb{C}_D \times \mathbb{R} \mapsto \mathbb{R}$  defined by (3.1) is continuous and maps bounded sets into bounded sets.*<sup>1</sup>

*Proof.* (i) Assume that  $\mathcal{A} \subset \mathbb{C}_D \times \mathbb{R}$  and  $\gamma \in (0, \infty)$  are such that  $\|\ell\|_\infty + |d| \leq \gamma$  for each  $(\ell, d) \in \mathcal{A}$  and that there is a sequence  $\{a(\ell_n, d_n)\}_{n=1}^\infty \subset a(\mathcal{A})$  such that  $\lim_{n \rightarrow \infty} a(\ell_n, d_n) = \infty$  or  $\lim_{n \rightarrow \infty} a(\ell_n, d_n) = -\infty$ . Let the former possibility occur. Then, by (3.1), we have  $0 = \lim_{n \rightarrow \infty} \Psi_{\ell_n, d_n}(a(\ell_n, d_n)) \geq \lim_{n \rightarrow \infty} (-\gamma + T\phi^{-1}(a(\ell_n, d_n) - \gamma)) = \infty$ , a contradiction. The latter possibility can be argued similarly.

(ii) Let  $\lim_{n \rightarrow \infty} (\ell_n, d_n) = (\ell_0, d_0)$  in  $\mathbb{C}_D \times \mathbb{R}$ . By (i) the sequence  $\{a(\ell_n, d_n)\}_{n=1}^\infty$  is bounded and hence we can choose a subsequence such that  $\lim_{n \rightarrow \infty} a(\ell_{k_n}, d_{k_n}) = a_0 \in \mathbb{R}$ . By (3.1), we get

$$0 = \Psi_{\ell_{k_n}, d_{k_n}}(a(\ell_{k_n}, d_{k_n})) = d_{k_n} + \int_0^T \phi^{-1}(a(\ell_{k_n}, d_{k_n}) + \ell_{k_n}(t)) \, dt,$$

which, for  $n \rightarrow \infty$ , yields

$$0 = d_0 + \int_0^T \phi^{-1}(a_0 + \ell_0(t)) \, dt.$$

Thus, with respect to Lemma 3.1, we have  $a_0 = a(\ell_0, d_0) = \lim_{n \rightarrow \infty} a(\ell_n, d_n)$ .  $\square$

**3.3 Lemma.** *The operator  $\mathcal{N} : \mathbb{C}_D^1 \mapsto \mathbb{C}_D$  given by*

$$(3.2) \quad (\mathcal{N}(x))(t) = \int_0^t f(s, x(s), x'(s)) \, ds + \sum_{i=1}^m [\phi(M_i(x'(t_i))) - \phi(x'(t_i))] \chi_{(t_i, T]}(t)$$

*is absolutely continuous.*<sup>2</sup>

*Proof.* The continuity of  $\mathcal{N}$  follows from the continuity of all the mappings involved in the right-hand side of (3.2). Furthermore, let  $\mathcal{H} \subset \mathbb{C}_D^1$  be bounded. We need to show that the closure  $\overline{\mathcal{N}(\mathcal{H})}$  of  $\mathcal{N}(\mathcal{H})$  in  $\mathbb{C}_D$  is compact. To this aim, let  $\|x\|_D \leq \gamma < \infty$  for each  $x \in \mathcal{H}$ . Then there are  $c \in (0, \infty)$  and  $h \in \mathbb{L}_1$  such that

$$\sum_{i=1}^m [\phi(M_i(x'(t_i))) - \phi(x'(t_i))] \leq c \quad \text{and} \quad |f(t, x(t), x'(t))| \leq h(t) \quad \text{a.e. on } [0, T]$$

for all  $x \in \mathcal{H}$ . Therefore

$$(3.3) \quad \|\mathcal{N}(x)\|_\infty \leq \|h\|_1 + c \quad \text{for each } x \in \mathcal{H}.$$

<sup>1</sup>The norm of  $(\ell, d) \in \mathbb{C}_D \times \mathbb{R}$  is defined by  $\|\ell\|_\infty + |d|$ .

<sup>2</sup>As usual,  $\chi_M$  stands for the characteristic function of the set  $M \subset \mathbb{R}$ .

Put  $(\mathcal{N}_1(x))(t) = \int_0^t f(s, x(s), x'(s)) \, ds$ . Then, for  $t_1, t_2 \in [0, T]$ , we have

$$|(\mathcal{N}_1(x))(t_2) - (\mathcal{N}_1(x))(t_1)| \leq \left| \int_{t_1}^{t_2} h(s) \, ds \right|,$$

wherefrom, by (3.3), we deduce that the functions in  $\mathcal{N}_1(\mathcal{H})$  are uniformly bounded and equicontinuous on  $[0, T]$ . Hence, making use of the Arzelà-Ascoli Theorem in  $\mathbb{C}$  (the space of functions continuous on  $[0, T]$  with the norm  $\|\cdot\|_\infty$ ), we get that each sequence in  $\mathcal{N}_1(\mathcal{H})$  contains a subsequence convergent with respect to the norm  $\|\cdot\|_\infty$ . This shows that  $\overline{\mathcal{N}_1(\mathcal{H})}$  is compact in  $\mathbb{C}_D$ . We know that the operator  $\mathcal{N}_2 = \mathcal{N} - \mathcal{N}_1$  is continuous. By (3.3), it maps bounded sets into bounded sets. Moreover, its values are contained in an  $m$ -dimensional subspace of  $\mathbb{C}_D$ . Thus,  $\overline{\mathcal{N}_2(\mathcal{H})}$  is compact in  $\mathbb{C}_D$ .  $\square$

**3.4 Theorem.** *Let  $a : \mathbb{C}_D \times \mathbb{R} \mapsto \mathbb{R}$  and  $\mathcal{N} : \mathbb{C}_D^1 \mapsto \mathbb{C}_D$  be respectively defined by (3.1) and (3.2). Furthermore define  $\mathcal{J} : \mathbb{C}_D^1 \mapsto \mathbb{C}_D^1$  by*

$$(3.4) \quad (\mathcal{J}(x))(t) = \sum_{i=1}^m [J_i(x(t_i)) - x(t_i)] \chi_{(t_i, T]}(t)$$

and

$$(3.5) \quad (\mathcal{F}(x))(t) = \int_0^t \phi^{-1} \left( a(\mathcal{N}(x), (\mathcal{J}(x))(T)) + (\mathcal{N}(x))(s) \right) \, ds \\ + x(0) + x'(0) - x'(T) + (\mathcal{J}(x))(t).$$

Then  $\mathcal{F} : \mathbb{C}_D^1 \mapsto \mathbb{C}_D^1$  is an absolutely continuous operator. Moreover,  $u$  is a solution of the problem (1.1) – (1.3) if and only if  $\mathcal{F}(u) = u$ .

*Proof.* For  $x \in \mathbb{C}_D^1$  and  $t \in [0, T]$ , we have

$$(3.6) \quad (\mathcal{F}(x))'(t) = \phi^{-1} \left( a(\mathcal{N}(x), (\mathcal{J}(x))(T)) + (\mathcal{N}(x))(t) \right).$$

Since the mappings  $a$ ,  $\mathcal{N}$  and  $\mathcal{J}$  included in (3.5) and (3.6) are continuous, it follows that  $\mathcal{F}$  is continuous in  $\mathbb{C}_D^1$ .

Choose an arbitrary bounded set  $\mathcal{H} \subset \mathbb{C}_D^1$ . We will show that then the set  $\overline{\mathcal{F}(\mathcal{H})}$  is compact in  $\mathbb{C}_D^1$ . Let a sequence  $\{v_n\} \subset \mathcal{F}(\mathcal{H})$  be given. It suffices to show that it contains a subsequence convergent in  $\mathbb{C}_D^1$ . Let  $\{x_n\} \subset \mathcal{H}$  be such that  $v_n = \mathcal{F}(x_n)$  for  $n \in \mathbb{N}$ . By Lemma 3.3, there is a subsequence  $\{x_{k_n}\}$  such that  $\{\mathcal{N}(x_{k_n})\}$  is convergent in  $\mathbb{C}_D$ . According to (3.3) and (3.4), there exists  $\gamma \in (0, \infty)$  such that  $\|\mathcal{N}(x)\|_\infty + |(\mathcal{J}(x))(T)| \leq \gamma$  for all  $x \in \mathcal{H}$ . Hence, by Lemma 3.2, the sequence  $\{a(\mathcal{N}(x_{k_n}), (\mathcal{J}(x_{k_n}))(T))\} \subset \mathbb{R}$  is bounded and we can choose a subsequence  $\{x_{\ell_n}\} \subset \{x_{k_n}\}$  such that  $\{a(\mathcal{N}(x_{\ell_n}), (\mathcal{J}(x_{\ell_n}))(T)) + \mathcal{N}(x_{\ell_n})\}$  is convergent in  $\mathbb{C}_D$ . Consequently,  $\{(\mathcal{F}(x_{\ell_n}))'\}$  and  $\{\mathcal{F}(x_{\ell_n})\}$  are convergent in  $\mathbb{C}_D$ , as well. Finally, by a direct computation we check that (1.1)–(1.3) is equivalent to the problem  $u = \mathcal{F}(u)$ . For more details, see our preprint [15].  $\square$

## 4. Proofs of the main results

**Sketch of the proof of Theorem 2.1.** We can modify the arguments and constructions of [13], where the case  $\phi(y) \equiv y$  is considered. By virtue of Theorem 3.4, the problem (1.1)–(1.3) has a solution if and only if the operator  $\mathcal{F}$  which is defined by (3.5) has a fixed point. To prove it we argue as follows: (i) we construct an auxiliary operator  $\tilde{\mathcal{F}}$  and prove that its Leray-Schauder topological degree is nonzero and consequently  $\tilde{\mathcal{F}}$  has a fixed point  $u$ ; (ii) using the method of a priori estimates we show that  $u$  is a fixed point of  $\mathcal{F}$  satisfying (2.1). Since the realization of these ideas is quite close to the arguments of [13], we skip it. Detailed computation can be found in our preprint [15].

**Proof of Theorem 2.2.** STEP 1. Define

$$(4.1) \quad \beta_j(y) = \begin{cases} c_j & \text{for } y < c_j, \\ y & \text{for } c_j \leq y \leq d_j, \\ d_j & \text{for } y > d_j \end{cases} \quad j = 1, 2, \dots, m+1;$$

$$(4.2) \quad \tilde{f}(t, x, y) = f(t, x, \beta_j(y)) + \frac{y - \beta_j(y)}{|y - \beta_j(y)| + 1}$$

for a.e.  $t \in (t_{j-1}, t_j]$ ,  $x, y \in \mathbb{R}$ ,  $j = 1, 2, \dots, m+1$ ;

and

$$(4.3) \quad \tilde{M}_i(y) = M_i(\beta_i(y)) + \frac{y - \beta_i(y)}{|y - \beta_i(y)| + 1} \quad \text{for } y \in \mathbb{R}, \quad i = 1, 2, \dots, m.$$

Now, consider the auxiliary problem

$$(4.4) \quad (\phi(u'(t)))' = \tilde{f}(t, u(t), u'(t)) \quad \text{a.e. on } [0, T];$$

$$(4.5) \quad u(t_i+) = J_i(u(t_i)), \quad u'(t_i+) = \tilde{M}_i(u'(t_i)), \quad i = 1, 2, \dots, m,$$

$$(4.6) \quad u(0) = u(T), \quad \beta_1(u'(0)) = u'(T).$$

We see that  $\tilde{f}$  and  $\tilde{M}_i$  have the same properties as  $f$  and  $M_i$ . In particular,  $\tilde{f}$  satisfies (1.11) with  $\omega(s) \equiv 1$ ,  $\tilde{M}_i$  fulfils (1.9) and  $\sigma_1/\sigma_2$  are lower/upper functions for (4.4)–(4.6). Since we work with (4.6) instead of (1.3), we have to replace the expression  $x(0) + x'(0) - x'(T)$  in (3.5) by  $x(0) + \beta_1(x'(0)) - x'(T)$ . Then we get the existence of a solution  $u$  of (4.4)–(4.6) satisfying (2.1) in the same way as in the proof of Theorem 2.1 for (1.1)–(1.3).

STEP 2. Having the solution  $u$  of (4.4)–(4.6), it remains to show that (2.2) is true.

(i) Let  $j \in \{1, 2, \dots, m+1\}$  and  $\xi \in [t_{j-1}, t_j)$  be such that

$$(4.7) \quad \sup\{u'(t) : t \in [0, T]\} = u'(\xi+) > d_j.$$

Then there is  $\delta > 0$  such that  $(\xi, \xi + \delta) \subset (t_{j-1}, t_j)$  and  $u' > d_j$  on  $(\xi, \xi + \delta)$ . By (1.12),

$$(\phi(u'(t)))' = f(t, u(t), d_j) + \frac{u'(t) - d_j}{u'(t) - d_j + 1} > 0 \quad \text{for a.e. } t \in (\xi, \xi + \delta),$$

i.e.  $\phi(u'(t)) > \phi(u'(\xi+))$  and so  $u'(t) > u'(\xi+)$  for each  $t \in (\xi, \xi + \delta)$ , which contradicts (4.7).

(ii) Assume that

$$(4.8) \quad \sup\{u'(t) : t \in [0, T]\} = u'(t_j) > d_j \quad \text{for some } t_j \in D.$$

If  $j = m + 1$ , i.e.  $u'(T) > d_{m+1}$ , then, by (1.12), we have also  $u'(T) > d_1$ . Since (4.1) and (4.6) imply  $u'(T) \leq d_1$ , we get a contradiction.

If  $j < m + 1$ , then

$$\widetilde{M}_j(u'(t_j)) = M_j(d_j) + \frac{u'(t) - d_j}{u'(t) - d_j + 1} > M_j(d_j) \geq d_{j+1},$$

so  $u'(t_j+) > d_{j+1}$ . By part (i) we know that  $u'(t) - d_{j+1}$  cannot achieve a positive maximum inside  $(t_j, t_{j+1})$ . Consequently, we have  $u'(t_{j+1}) > d_{j+1}$ . Repeating this procedure we get  $u'(T) > d_{m+1}$  and a contradiction as before.

We have proved that  $u'(t) \leq d_j$  on  $(t_{j-1}, t_j]$ ,  $j = 1, 2, \dots, m + 1$ . The remaining inequalities in (2.2) can be derived analogously. Finally, since  $u$  fulfils (2.2),  $u$  is a solution of (1.1)–(1.3).

**Sketch of the proof of Theorem 2.3.** We borrow ideas of [14], where non-ordered lower/upper functions to periodic impulsive problem without  $\phi$ -Laplacian ( $\phi(y) = y$ ) have been studied. Here, we define the operator  $\mathcal{F}$  by (3.5). Then, according to  $\mathcal{F}$ , we construct auxiliary operators and compute their Leray-Schauder degrees by a similar procedure as in [14]. For this we need a priori estimates of solutions of corresponding auxiliary problems. Now we consider problems with  $\phi$ -Laplacians but the basic evaluation of estimates of  $\phi(u')$  are similar to those of  $u'$  in [14] and hence we omit their computation here. For details see our preprint [16].

**Proof of Theorem 2.4.** First, we will prove the following a priori estimate:

CLAIM. *There exist  $a_j \in (0, \infty)$ ,  $j = 1, 2, \dots, m + 1$ , such that for each function  $u \in \mathbb{C}_D^1$  satisfying (1.2), (1.3), (2.2) and (2.3), the estimates*

$$(4.9) \quad |u(t)| \leq a_j \quad \text{for } t \in (t_{j-1}, t_j], \quad j = 1, 2, \dots, m + 1$$

are valid.



Indeed, let  $u$  satisfy the assumptions of CLAIM and let

$$\rho_0 = \max\{\|\sigma_1\|_\infty, \|\sigma_2\|_\infty\} \quad \text{and} \quad \gamma_i = \max\{|c_i|, |d_i|\}, \quad i = 1, 2, \dots, m + 1.$$

(i) If  $t_u \in [0, t_1]$ , then  $|u(t)| \leq \gamma_1 t_1 + \rho_0$  for  $t \in [0, t_1]$ . Put  $a_1^0 = \gamma_1 t_1 + \rho_0$  and  $b_1^0 = \max\{|J_1(x)| : x \in [-a_1^0, a_1^0]\}$ . Then  $|u(t)| \leq \gamma_2 (t_2 - t_1) + b_1^0$  for  $t \in (t_1, t_2]$ . Further, put  $a_2^0 = \gamma_2 (t_2 - t_1) + b_1^0$  and  $b_2^0 = \max\{|J_2(x)| : x \in [-a_2^0, a_2^0]\}$ . Then  $|u(t)| \leq \gamma_3 (t_3 - t_2) + b_2^0$  for  $t \in (t_2, t_3]$ . By induction we get that  $|u(t)| \leq a_i^0$  for  $t \in (t_{i-1}, t_i]$ , where  $a_{i+1}^0 = \gamma_{i+1} (t_{i+1} - t_i) + \max\{|J_i(x)| : x \in [-a_i^0, a_i^0]\}$ ,  $i = 1, 2, \dots, m$ .  
(ii) If  $t_u \in (t_j, t_{j+1}]$  for some  $j \in \{1, 2, \dots, m\}$ , we get similarly as in (i) that  $|u(t)| \leq a_i^j$  for  $t \in (t_{i-1}, t_i]$ ,  $i = 1, 2, \dots, m + 1$ , where  $a_{j+1}^j = \gamma_{j+1} (t_{j+1} - t_j) + \rho_0$ ,  $a_{i+1}^j = \gamma_{i+1} (t_{i+1} - t_i) + \max\{|J_i(x)| : x \in [-a_i^j, a_i^j]\}$ ,  $i = 1, 2, \dots, j - 1, j + 1, \dots, m$ ,  $a_1^j = \gamma_1 t_1 + a_{m+1}^j$ .

Setting

$$a_j = \max\{\rho_0, a_j^0, a_j^1, \dots, a_j^m\} \quad \text{for } j = 1, 2, \dots, m + 1,$$

we complete the proof of CLAIM.

Now, take  $\beta_j$  by (4.1) and for  $a_j$  of CLAIM put

$$\alpha_j(x) = \begin{cases} -a_j & \text{for } x < -a_j, \\ x & \text{for } -a_j \leq x \leq a_j, \\ a_j & \text{for } x > a_j \end{cases}$$

and

$$\begin{aligned} \tilde{f}(t, x, y) &= f(t, \alpha_j(x), \beta_j(y)) + \frac{y - \beta_j(y)}{|y - \beta_j(y)| + 1} \\ &\text{for a.e. } t \in (t_{j-1}, t_j], \text{ all } x, y \in \mathbb{R}, \quad j = 1, 2, \dots, m + 1. \end{aligned}$$

Finally, define  $\widetilde{M}_i$  by (4.3). We see that all assumptions of Theorem 2.3 are satisfied for the problem (4.4)–(4.6) and consequently it has a solution  $u$  satisfying (2.3). As in the proof of Theorem 2.2, Step 2, we get that  $u$  fulfils (2.2). Hence  $u$  satisfies (1.2), (1.3) and, by CLAIM, also (4.8). Therefore,  $u$  is a solution of (1.1)–(1.3).

## References

- [1] A. CABADA AND R. POUSO. Existence result for the problem  $(\phi(u'))' = f(t, u, u')$  with periodic and Neumann boundary conditions. *Nonlinear Anal., Theory Methods Appl.* **30** (1997), 1733–1742.
- [2] A. CABADA, P. HABETS AND R. POUSO. Lower and upper solutions for the periodic problem associated with a  $\phi$ -Laplacian equation. *EQUADIFF 1999 - International Conference on Differential Equations*, Vol. 1, 2 (Berlin, 1999), 491–493, World Sci. Publishing, River Edge, NJ, 2000.
- [3] M. CHERPION, C. DE COSTER AND P. HABETS. Monotone iterative methods for boundary value problems. *Differ. Integral. Equ.* **12** (3) (1999), 309–338.

- [4] M. DEL PINO, R. MANÁSEVICH AND A. MURÚA. Existence and multiplicity of solutions with prescribed period for a second order quasilinear O.D.E. *Nonlinear Anal., Theory Methods Appl.* **18** (1992), 79–92.
- [5] C. FABRY AND D. FAYYAD. Periodic solutions of second order differential equation with a  $p$ -Laplacian and asymmetric nonlinearities. *Rend. Ist. Mat. Univ. Trieste* **24** (1992), 207–227.
- [6] P. GIRG. Neumann and periodic boundary-value problems for quasilinear ordinary differential equations with a nonlinearity in the derivative. *Electron. J. Differ. Equ.*, 2000, Paper No.63, 28 p., electronic only (2000).
- [7] P. JEBELEAN AND J. MAWHIN. Periodic solutions of singular nonlinear perturbations of the ordinary  $p$ -Laplacian. *Adv. Nonlinear Stud.* **2** (2002), 299–312.
- [8] BIN LIU. Multiplicity results for periodic solutions of a second order quasilinear ordinary differential equations with asymmetric nonlinearities. *Nonlinear Anal., Theory Methods Appl.* **33** (2) (1998), 139–160.
- [9] BING LIU. Periodic solutions of dissipative dynamical systems with singular potential and  $p$ -Laplacian. *Ann. Pol. Math.* **79** (2)(2002), 109–120.
- [10] R. MANÁSEVICH AND J. MAWHIN. Periodic solutions for nonlinear systems with  $p$ -Laplacian like operators. *J. Differ. Equations* **145** (1998), 367–393.
- [11] J. MAWHIN. Some boundary value problems for Hartman-type perturbations of the ordinary vector  $p$ -Laplacian. *Nonlinear Anal., Theory Methods Appl.* **40** (2000), 497–503.
- [12] J. MAWHIN AND A. UREÑA. A Hartman-Nagumo inequality for the vector ordinary  $p$ -Laplacian and application to nonlinear boundary value problem. *J. Inequal. Appl.* **7**(5) (2002), 701–725.
- [13] I. RACHŮNKOVÁ AND M. TVRĎÝ. Nonmonotone impulse effects in second order periodic boundary value problems. *Abstr. Anal. Appl.*, to appear. [Preprint entitled *Periodic boundary value problems for nonlinear second order differential equations with impulses - Part I* is available as <http://www.math.cas.cz/~tvrdy/i2.ps>.]
- [14] I. RACHŮNKOVÁ AND M. TVRĎÝ. Existence results for impulsive second order periodic problems, submitted. [Preprint entitled *Periodic boundary value problems for nonlinear second order differential equations with impulses - Part III* is available as <http://www.math.cas.cz/~tvrdy/i4.ps>.]
- [15] I. RACHŮNKOVÁ AND M. TVRĎÝ. Second Order Periodic Problem with  $\phi$ -Laplacian and Impulses - Part I. [Preprint is available as <http://www.math.cas.cz/~tvrdy/lap11.ps>.]
- [16] I. RACHŮNKOVÁ AND M. TVRĎÝ. Second Order Periodic Problem with  $\phi$ -Laplacian and Impulses - Part II. [Preprint is available as <http://www.math.cas.cz/~tvrdy/lap12.ps>.]
- [17] S. STANĚK. Periodic boundary value problem for second order functional differential equations. *Math. Notes, Miskolc* **1** (2000), 63–81.
- [18] S. STANĚK. On solvability of singular periodic boundary value problems. *Nonlinear Oscil.* **4** (2001), 529–538.
- [19] PING YAN. Nonresonance for one-dimensional  $p$ -Laplacian with regular restoring. *J. Math. Anal. Appl.* **285** (2003), 141–154.

Irena Rachůnková, Department of Mathematics, Palacký University, 779 00 OLOMOUC, Tomkova 40, Czech Republic (e-mail: [rachunko@risc.upol.cz](mailto:rachunko@risc.upol.cz))

Milan Tvrďý, Mathematical Institute, Academy of Sciences of the Czech Republic, 115 67 PRAHA 1, Žitná 25, Czech Republic (e-mail: [tvrdy@math.cas.cz](mailto:tvrdy@math.cas.cz))