

Asymptotic formula for oscillatory solutions of some singular nonlinear differential equation*

IRENA RACHŮNKOVÁ, LUKÁŠ RACHŮNEK

Department of Mathematics,
Faculty of Science, Palacký University,
17. listopadu 12, 771 46 Olomouc, Czech Republic
e-mail: rachunko@inf.upol.cz, rachunekl@phoenix.inf.upol.cz

Abstract

Singular differential equation $(p(t)u')' = p(t)f(u)$ is investigated. Here f is Lipschitz continuous on \mathbb{R} and has at least two zeros 0 and $L > 0$. The function p is continuous on $[0, \infty)$, has a positive continuous derivative on $(0, \infty)$ and $p(0) = 0$. An asymptotic formula for oscillatory solutions is derived.

Mathematics Subject Classification 2010: 34D05, 34A12, 34B40

Key words: Singular ordinary differential equation of the second order, time singularities, asymptotic formula, oscillatory solutions.

1 Introduction

In this paper we investigate the equation

$$(p(t)u')' = p(t)f(u), \quad t \in (0, \infty), \quad (1)$$

where f satisfies

$$f \in Lip_{loc}(\mathbb{R}), \quad f(0) = f(L) = 0, \quad f(x) < 0, \quad x \in (0, L), \quad (2)$$

$$\exists \bar{B} \in (-\infty, 0) : f(x) > 0, \quad x \in [\bar{B}, 0), \quad (3)$$

$$F(\bar{B}) = F(L), \quad \text{where } F(x) = - \int_0^x f(z) dz, \quad x \in \mathbb{R}, \quad (4)$$

*Supported by the Council of Czech Government MSM 6198959214

and p fulfils

$$p \in C[0, \infty) \cap C^1(0, \infty), \quad p(0) = 0, \quad (5)$$

$$p'(t) > 0, \quad t \in (0, \infty), \quad \lim_{t \rightarrow \infty} \frac{p'(t)}{p(t)} = 0. \quad (6)$$

Equation (1) is a generalization of the equation

$$u'' + \frac{k-1}{t}u' = f(u), \quad t \in (0, \infty), \quad (7)$$

which arises for $k > 1$ and special forms of f in many areas. For example: In the study of phase transitions of Van der Waals fluids [3], [10], [23], in population genetics, where it serves as a model for the spatial distribution of the genetic composition of a population [8], [9], in the homogeneous nucleation theory [1], in relativistic cosmology for description of particles which can be treated as domains in the universe [17], in the nonlinear field theory, in particular, when describing bubbles generated by scalar fields of the Higgs type in the Minkowski spaces [7]. Numerical simulations of solutions of (1), where f is a polynomial with three zeros have been presented in [6], [14], [16]. Close problems about the existence of positive solutions can be found in [2], [4], [5].

Due to $p(0) = 0$, equation (1) has a singularity at $t = 0$.

Definition 1 A function $u \in C^1[0, \infty) \cap C^2(0, \infty)$ which satisfies equation (1) for all $t \in (0, \infty)$ is called a *solution* of (1).

Definition 2 Let u be a solution of equation (1) and let L be of (2). Denote $u_{\text{sup}} = \sup\{u(t) : t \in [0, \infty)\}$. If $u_{\text{sup}} < L$ ($u_{\text{sup}} = L$ or $u_{\text{sup}} > L$), then u is called a *damped solution* (a *bounding homoclinic solution* or an *escape solution*).

These three types of solutions have been investigated in [18]-[22]. In particular, the existence of damped oscillatory solutions which converge to 0 has been proved in [22].

The main result of this paper is contained in Section 3 in Theorem 8, where we provide an asymptotic formula for damped oscillatory solutions of equation (1).

2 Existence of oscillatory solutions

Here we will study solutions of (1) satisfying the initial conditions

$$u(0) = B, \quad u'(0) = 0, \quad (8)$$

with a parameter $B \leq L$. Reason is that we focus our attention on damped solutions of (1) and that each solution u of (1) must fulfil $u'(0) = 0$ (see [22]).

First, we bring two theorems about the existence of damped and oscillatory solutions.

Theorem 3 [22] Assume that (2)-(6) hold. Then for each $B \in [\bar{B}, L)$ problem (1), (8) has a unique solution. This solution is damped.

Theorem 4 Assume that (2)-(6) hold. Further, let there exists $k_0 \in (0, \infty)$ such that

$$p \in C^2(0, \infty), \quad \limsup_{t \rightarrow \infty} \left| \frac{p''(t)}{p'(t)} \right| < \infty, \quad \liminf_{t \rightarrow \infty} \frac{p(t)}{t^{k_0}} \in (0, \infty], \quad (9)$$

$$\lim_{x \rightarrow 0^+} \frac{f(x)}{x} < 0, \quad \lim_{x \rightarrow 0^-} \frac{f(x)}{x} < 0. \quad (10)$$

Then for each $B \in [\bar{B}, L)$ problem (1), (8) has a unique solution u . If $B \neq 0$, then the solution u is damped and oscillatory with decreasing amplitudes and

$$\lim_{t \rightarrow \infty} u(t) = 0. \quad (11)$$

Proof. The assertion follows from Theorems 5,11 and 17 in [22]. \square

Example 5 The functions

- $p(t) = t^k, \quad p(t) = t^k \ln(t^\ell + 1), \quad k, \ell \in (0, \infty),$
- $p(t) = t + \alpha \sin t, \quad \alpha \in (-1, 1),$
- $p(t) = \frac{t^k}{1+t^\ell}, \quad k, \ell \in (0, \infty), \ell < k,$

satisfy (5), (6) and (9).

The functions

- $p(t) = \ln(t+1), \quad p(t) = \arctan t, \quad p(t) = \frac{t^k}{1+t^k}, \quad k \in (0, \infty),$

satisfy (5), (6), but not (9) (the third condition).

The function

- $p(t) = t^k + \alpha \sin t^k, \quad \alpha \in (-1, 1), k \in (1, \infty),$

satisfy (5), (6) but not (9) (the second and third conditions).

Example 6 Let $k \in (0, \infty)$.

- The function

$$f(x) = \begin{cases} -kx & \text{for } x \leq 0 \\ x(x-1) & \text{for } x > 0, \end{cases}$$

satisfies (2) with $L = 1$, (3), (4) with $\bar{B} = -(3k)^{-1/2}$ and (10).

- The function

$$f(x) = \begin{cases} kx^2 & \text{for } x \leq 0 \\ x(x-1) & \text{for } x > 0, \end{cases}$$

satisfies (2) with $L = 1$, (3), (4) with $\bar{B} = -(2k)^{-1/3}$ but not (10) (the second condition).

In the next section, the generalized Matell's theorem which can be found as Theorem 6.5 in the monograph by I. Kiguradze will be useful. For our purpose we provide its following special case.

Consider an interval $J \subset \mathbb{R}$. We write $AC(J)$ for the set of functions absolutely continuous on J and $AC_{loc}(J)$ for the set of functions belonging to $AC(I)$ for each compact interval $I \subset J$. Choose $t_0 > 0$ and a function matrix $A(t) = (a_{i,j}(t))_{i,j \leq 2}$ which is defined on (t_0, ∞) . Denote by $\lambda(t)$ and $\mu(t)$ eigenvalues of $A(t)$, $t \in (t_0, \infty)$. Further, suppose

$$\lambda = \lim_{t \rightarrow \infty} \lambda(t) \quad \text{and} \quad \mu = \lim_{t \rightarrow \infty} \mu(t)$$

be different eigenvalues of the matrix $A = \lim_{t \rightarrow \infty} A(t)$ and let \mathbf{l} and \mathbf{m} be eigenvectors of A corresponding to λ and μ , respectively.

Theorem 7 [11] *Assume that*

$$a_{i,j} \in AC_{loc}(t_0, \infty), \quad \left| \int_{t_0}^{\infty} a'_{i,j}(t) dt \right| < \infty, \quad i, j = 1, 2, \quad (12)$$

and that there exists $c_0 > 0$ such that

$$\int_s^t \operatorname{Re}(\lambda(\tau) - \mu(\tau)) d\tau \leq c_0, \quad t_0 \leq s < t, \quad (13)$$

or

$$\int_{t_0}^{\infty} \operatorname{Re}(\lambda(\tau) - \mu(\tau)) d\tau = \infty, \quad \int_s^t \operatorname{Re}(\lambda(\tau) - \mu(\tau)) d\tau \geq -c_0, \quad t_0 \leq s < t. \quad (14)$$

Then the differential system

$$\mathbf{x}'(t) = A(t)\mathbf{x}(t) \quad (15)$$

has a fundamental system of solutions $\mathbf{x}(t)$, $\mathbf{y}(t)$ such that

$$\lim_{t \rightarrow \infty} \mathbf{x}(t) e^{-\int_{t_0}^t \lambda(\tau) d\tau} = \mathbf{l}, \quad \lim_{t \rightarrow \infty} \mathbf{y}(t) e^{-\int_{t_0}^t \mu(\tau) d\tau} = \mathbf{m}. \quad (16)$$

3 Asymptotic formula

In order to derive an asymptotic formula for a damped oscillatory solution u of problem (1), (8), we need a little stronger assumption than (10). In particular, the function $\frac{f(x)}{x}$ should have a negative derivative at $x = 0$.

Theorem 8 *Assume that (2)-(6) and (9) hold. Assume moreover that there exist $\eta > 0$ and $c > 0$ such that*

$$\frac{f(x)}{x} \in AC[-\eta, \eta], \quad \lim_{x \rightarrow 0} \frac{f(x)}{x} = -c. \quad (17)$$

Then for each $B \in [\bar{B}, L)$ problem (1), (8) has a unique solution u . If $B \neq 0$, then the solution u is damped and oscillatory with decreasing amplitudes such that

$$\limsup_{t \rightarrow \infty} \sqrt{p(t)}|u(t)| < \infty. \quad (18)$$

Proof. Step 1. *Construction of an auxiliary linear differential system.* Choose $B \in [\bar{B}, L)$, $B \neq 0$. By Theorem 4, problem (1), (8) has a unique oscillatory solution u with decreasing amplitudes and satisfying (11). Having this solution u , define a linear differential equation

$$v'' + \frac{p'(t)}{p(t)}v' = \frac{f(u(t))}{u(t)}v, \quad (19)$$

and the corresponding linear differential system

$$x'_1 = x_2, \quad x'_2 = \frac{f(u(t))}{u(t)}x_1 - \frac{p'(t)}{p(t)}x_2. \quad (20)$$

Denote

$$A(t) = (a_{i,j}(t))_{i,j \leq 2} = \begin{pmatrix} 0 & 1 \\ \frac{f(u(t))}{u(t)} & -\frac{p'(t)}{p(t)} \end{pmatrix}, \quad A = \begin{pmatrix} 0 & 1 \\ -c & 0 \end{pmatrix}.$$

By (6), (11) and (17),

$$A = \lim_{t \rightarrow \infty} A(t).$$

Eigenvalues of A are numbers $\lambda = i\sqrt{c}$ and $\mu = -i\sqrt{c}$, eigenvectors of A are $\mathbf{l} = (1, i\sqrt{c})$ and $\mathbf{m} = (1, -i\sqrt{c})$, respectively. Denote

$$D(t) = \left(\frac{p'(t)}{2p(t)} \right)^2 + \frac{f(u(t))}{u(t)}, \quad t \in (0, \infty).$$

Then eigenvalues of $A(t)$ have the form

$$\lambda(t) = -\frac{p'(t)}{2p(t)} + \sqrt{D(t)}, \quad \mu(t) = -\frac{p'(t)}{2p(t)} - \sqrt{D(t)}, \quad t \in (0, \infty). \quad (21)$$

We see that

$$\lim_{t \rightarrow \infty} \lambda(t) = \lambda, \quad \lim_{t \rightarrow \infty} \mu(t) = \mu.$$

Step 2. *Verification of the assumptions of Theorem 7.* Due to (6), (11) and (17), we can find $t_0 > 0$ such that

$$u(t_0) \neq 0, \quad |u(t)| \leq \eta, \quad D(t) < 0, \quad t \in (t_0, \infty). \quad (22)$$

Therefore, by (17),

$$a_{21}(t) = \frac{f(u(t))}{u(t)} \in AC_{loc}(t_0, \infty),$$

and so

$$\left| \int_{t_0}^{\infty} \left(\frac{f(u(t))}{u(t)} \right)' dt \right| = \left| \lim_{t \rightarrow \infty} \frac{f(u(t))}{u(t)} - \frac{f(u(t_0))}{u(t_0)} \right| = \left| -c - \frac{f(u(t_0))}{u(t_0)} \right| < \infty.$$

Further, by (9), $a_{22}(t) = -\frac{p'(t)}{p(t)} \in C^1(t_0, \infty)$. Hence, due to (6),

$$\left| \int_{t_0}^{\infty} \left(\frac{p'(t)}{p(t)} \right)' dt \right| = \left| \lim_{t \rightarrow \infty} \frac{p'(t)}{p(t)} - \frac{p'(t_0)}{p(t_0)} \right| = \frac{p'(t_0)}{p(t_0)} < \infty.$$

Since $a_{11}(t) \equiv 0$ and $a_{12}(t) \equiv 1$, we see that (12) is satisfied. Using (21) we get $Re(\lambda(t) - \mu(t)) \equiv 0$. This yields

$$\int_s^t Re(\lambda(\tau) - \mu(\tau)) d\tau = 0 < c_0, \quad t_0 \leq s < t,$$

for any positive constant c_0 . Consequently (13) is valid.

Step 3. *Application of Theorem 7.* By Theorem 7 there exists a fundamental system $\mathbf{x}(t) = (x_1(t), x_2(t)), \mathbf{y}(t) = (y_1(t), y_2(t))$ of solutions of (20) such that (16) is valid. Hence

$$\lim_{t \rightarrow \infty} x_1(t) e^{-\int_{t_0}^t \lambda(\tau) d\tau} = 1, \quad \lim_{t \rightarrow \infty} y_1(t) e^{-\int_{t_0}^t \mu(\tau) d\tau} = 1. \quad (23)$$

Using (21) and (22) we get

$$\begin{aligned} \exp\left(-\int_{t_0}^t \lambda(\tau) d\tau\right) &= \exp\left(\int_{t_0}^t \left(\frac{p'(\tau)}{2p(\tau)} - \sqrt{D(\tau)}\right) d\tau\right) \\ &= \exp\left(\frac{1}{2} \ln \frac{p(t)}{p(t_0)}\right) \exp\left(-i \int_{t_0}^t \sqrt{|D(\tau)|} d\tau\right), \end{aligned}$$

and hence

$$\left| e^{-\int_{t_0}^t \lambda(\tau) d\tau} \right| = \sqrt{\frac{p(t)}{p(t_0)}}, \quad t \in (t_0, \infty).$$

Similarly

$$\left| e^{-\int_{t_0}^t \mu(\tau) d\tau} \right| = \sqrt{\frac{p(t)}{p(t_0)}}, \quad t \in (t_0, \infty).$$

Therefore (23) implies

$$1 = \lim_{t \rightarrow \infty} \left| x_1(t) e^{-\int_{t_0}^t \lambda(\tau) d\tau} \right| = \lim_{t \rightarrow \infty} |x_1(t)| \sqrt{\frac{p(t)}{p(t_0)}},$$

and

$$1 = \lim_{t \rightarrow \infty} \left| y_1(t) e^{-\int_{t_0}^t \mu(\tau) d\tau} \right| = \lim_{t \rightarrow \infty} |y_1(t)| \sqrt{\frac{p(t)}{p(t_0)}}.$$

Step 4. *Asymptotic formula.* In Step 1 we have assumed that u is a solution of equation (1), which means that

$$u''(t) + \frac{p'(t)}{p(t)}u'(t) = f(u(t)) \quad \text{for } t \in (0, \infty).$$

Consequently

$$u''(t) + \frac{p'(t)}{p(t)}u'(t) = \frac{f(u(t))}{u(t)}u(t) \quad \text{for } t \in (0, \infty),$$

and hence u is also a solution of equation (19). This yields that there are $c_1, c_2 \in \mathbb{R}$ such that $u(t) = c_1x_1(t) + c_2y_1(t)$, $t \in (0, \infty)$. Therefore

$$\limsup_{t \rightarrow \infty} \sqrt{p(t)}|u(t)| \leq (|c_1| + |c_2|)\sqrt{p(t_0)} < \infty.$$

□

Remark 9 Due to (9) and (18), we have for a solution u of Theorem 8

$$u(t) = O(t^{-k_0/2}) \quad \text{for } t \rightarrow \infty. \quad (24)$$

Example 10 Let $k \in (1, \infty)$.

- The functions $f(x) = x(x-1)$ and $f(x) = x(x-1)(x+2)$ satisfy all assumptions of Theorem 8.
- The functions $f(x) = x^{2k-1}(x-1)$ and $f(x) = x^{2k-1}(x-1)(x+2)$ satisfy (2)-(4) but not (17) (the second condition).

Example 11 Consider the initial problem

$$(t^2u')' = t^2u(u-5)(u+10), \quad u(0) = -3, \quad u'(0) = 0. \quad (25)$$

Here $L_0 = -10$, $L = 5$ and we can check that $\bar{B} < -3$. Further, all assumptions of Theorems 4 and 8 are fulfilled. Therefore, by Theorem 4, there exists a unique solution u of problem (25) which is damped and oscillatory and converges to 0. By Theorem 8, we have

$$\limsup_{t \rightarrow \infty} t|u(t)| < \infty, \quad \text{that is } u(t) = O\left(\frac{1}{t}\right) \quad \text{for } t \rightarrow \infty.$$

The behaviour of the solution $u(t)$ and of the function $tu(t)$ is presented on Figure 1.




Figure 1.

Remark 12 Our further research of this topic will be focused on a deeper investigation of all types of solutions defined in Definition 2. For example, we have proved in [18] and [22] that damped solutions of equation (1) can be either oscillatory or they have a finite number of zeros or no zero and converge to 0. A more precise characterization of behaviour of non-oscillatory solutions including their asymptotic formulas is an open problem. The same can be said about homoclinic solutions. In [20] we have found some conditions which guarantee their existence and we have shown that if u is a homoclinic solution of equation (1), then $\lim_{t \rightarrow \infty} u(t) = L$. In order to discover other existence conditions for homoclinic solutions we would like to estimate their convergence by proper asymptotic formulas.

Acknowledgement

The authors thank the referees for comments and suggestions.

References

- [1] F. F. Abraham, *Homogeneous Nucleation Theory*, Acad. Press, New York 1974.
- [2] H. Berestycki, P. L. Lions, L. A. Peletier, An ODE approach to the existence of positive solutions for semilinear problems in \mathbb{R}^N , *Indiana University Mathematics Journal* **30**, 1 (1981) 141–157.
- [3] V. Bongiorno, L. E. Scriven, H. T. Davis, Molecular theory of fluid interfaces, *J. Colloid and Interface Science* **57** (1967), 462–475.

- [4] D. Bonheure, J. M. Gomes, L. Sanchez, Positive solutions of a second-order singular ordinary differential equation, *Nonlinear Analysis* **61** (2005) 1383–1399.
- [5] M. Conti, L. Merizzi, S. Terracini, Radial solutions of superlinear equations in \mathbb{R}^N , Part I: A global variational approach, *Arch. Rational Mech. Anal.* **153** (2000) 291–316.
- [6] F. Dell’Isola, H. Gouin and G. Rotoli, Nucleation of spherical shell-like interfaces by second gradient theory: numerical simulations, *Eur. J. Mech B/Fluids* **15** (1996) 545–568.
- [7] G.H. Derrick, Comments on nonlinear wave equations as models for elementary particles, *J. Math. Physics* **5** (1965) 1252–1254.
- [8] P. C. Fife, *Mathematical aspects of reacting and diffusing systems*, Lecture notes in Biomathematics 28, Springer 1979.
- [9] R. A. Fischer, The wave of advance of advantageous genes, *Journ. of Eugenics* **7** (1937), 355–369.
- [10] H. Gouin and G. Rotoli, An analytical approximation of density profile and surface tension of microscopic bubbles for Van der Waals fluids, *Mech. Research Communic.* **24** (1997) 255–260.
- [11] I. Kiguradze, *Some Singular Boundary Value Problems for Ordinary Differential Equations*, ITU, Tbilisi 1975, 352 pages.
- [12] I. Kiguradze and T. Chanturia, *Asymptotic properties of solutions of nonautonomous ordinary differential equations*, Nauka, Moscow 1990, 429 pages.
- [13] I. Kiguradze and B. Shekhter, *Singular boundary value problems for ordinary differential equations of the second order*, Viniti, Moscow 1987, 97 pages.
- [14] G. Kitzhofer, O. Koch, P. Lima, E. Weinmüller, Efficient numerical solution of the density profile equation in hydrodynamics, *J. Sci. Comput.* **32**, 3 (2007) 411–424.
- [15] O. Koch, P. Kofler, E. Weinmüller, Initial value problems for systems of ordinary first and second order differential equations with a singularity of the first kind, *Analysis* **21** (2001) 373–389.
- [16] P.M. Lima, N.V.Chemetov, N.B. Konyukhova, A.I. Sukov, Analytical–numerical investigation of bubble-type solutions of nonlinear singular problems, *J. Comp. Appl. Math.* **189** (2006) 260–273.
- [17] A.P. Linde, *Particle Physics and Inflationary Cosmology*, Harwood Academic, Chur, Switzerland, 1990.

- [18] I. Rachůnková, J. Tomeček, Bubble-type solutions of nonlinear singular problem, *Mathematical and Computer Modelling* **51** (2010), 658–669.
- [19] I. Rachůnková, J. Tomeček, Strictly increasing solutions of a nonlinear singular differential equation arising in hydrodynamics, *Nonlinear Analysis* **72** (2010), 2114-2118.
- [20] I. Rachůnková, J. Tomeček, Homoclinic solutions of singular nonautonomous second order differential equations, *Boundary Value Problems*, Vol. 2009, Article ID 959636, 21 pages.
- [21] I. Rachůnková, J. Tomeček and J. Stryja, Oscillatory solutions of singular equations arising in hydrodynamics, *Advances in Difference Equations, Recent Trends in Differential and Difference Equations*, Vol. 2010, Article ID 872160, 13 pages.
- [22] I. Rachůnková, L. Rachůnek and J. Tomeček, Existence of oscillatory solutions of singular nonlinear differential equations, *Abstract and Applied Analysis*, to appear.
- [23] J. D. van der Waals, R. Kohnstamm, *Lehrbuch der Thermodynamik*, vol.1, Leipzig, 1908.